

Post-genomic approaches to understanding interactions between fungi and their environment

Ronald P. de Vries^{1,2}, Isabelle Benoit², Gunther Doehlemann³, Tetsuo Kobayashi⁴, Jon K. Magnuson⁵, Ellen A. Panisko⁵, Scott E. Baker⁵, and Marc-Henri Lebrun⁶

¹Fungal Physiology, CBS-KNAW Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands; corresponding author e-mail: r.devries@cbs.knaw.nl

²Microbiology & Kluiver Centre for Genomics of Industrial Fermentation, Utrecht University, Padualaan 8, 3584 C Utrecht, The Netherlands

³MPI for Terrestrial Microbiology, Department of Organismic Interactions, Karl von Frisch Str. 10, 35043 Marburg, Germany

⁴Department of Biological Mechanisms and Functions, Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-8601, Japan

⁵Fungal Biotechnology Team, Pacific Northwest National Laboratory, Richland, WA 99352, USA

⁶BIOGER-INRA, Grignon, Av Lucien Brétignières, 78 850 Thiverval, Grignon, France

Abstract: Fungi inhabit every natural and anthropogenic environment on Earth. They have highly varied life-styles including saprobes (using only dead biomass as a nutrient source), pathogens (feeding on living biomass), and symbionts (co-existing with other organisms). These distinctions are not absolute as many species employ several life styles (e.g. saprobe and opportunistic pathogen, saprobe and mycorrhiza). To efficiently survive in these different and often changing environments, fungi need to be able to modify their physiology and in some cases will even modify their local environment. Understanding the interaction between fungi and their environments has been a topic of study for many decades. However, recently these studies have reached a new dimension. The availability of fungal genomes and development of post-genomic technologies for fungi, such as transcriptomics, proteomics and metabolomics, have enabled more detailed studies into this topic resulting in new insights. Based on a Special Interest Group session held during IMC9, this paper provides examples of the recent advances in using (post-)genomic approaches to better understand fungal interactions with their environments.

Key words:

(post-)genomics
Aspergillus oryzae
Aspergillus niger
Phycomyces blakesleeianus
Thielavia terrestris
Ustilago maydis

Article info: Submitted 5 May 2011; Accepted 18 May 2011; Published 24 May 2011.

INTRODUCTION

The interaction between fungi and their environment is of major importance for saprobes, symbionts and pathogens and has been a topic of study for many decades. Currently, genome sequences are available for many fungi, including saprobes (Espagne *et al.* 2008, Galagan *et al.* 2003, 2005, Jeffries *et al.* 2007, Machida *et al.* 2005, Martinez *et al.* 2004, 2008, Ohm *et al.* 2010, Pel *et al.* 2007), plant pathogens (Cuomo *et al.* 2007, Dean *et al.* 2005, Kämpfer *et al.* 2006), human pathogens (Nierman *et al.* 2005) and mycorrhizae (Martin *et al.* 2008, 2010). Sequencing of additional fungal genomes is occurring at an increasing rate in many centres all over the world, but the two largest fungal sequencing programs are running in the US at the Joint Genome Institute of the Department of Energy (<http://genome.jgi-psf.org/programs/fungi/index.jsf>) and the Broad Institute (<http://www.broadinstitute.org/scientific-community/>

science/projects/fungal-genome-initiative/fungal-genome-initiative). These initiatives are providing genome resources for a representative subset of the fungi, enabling full-genome comparison of fungal biodiversity.

The availability of fungal genome sequences was followed by the development of post-genomic resources, of which transcriptomics was the first. Initially transcriptomics was mainly available for fungal species which were studied by large consortia of scientists, due to the high costs involved in developing micro-arrays. Most papers on fungal transcriptomics have therefore addressed species such as *Saccharomyces cerevisiae*, various *Aspergillus* species, *Candida albicans*, *Neurospora crassa* and *Magnaporthe grisea* (Andersen *et al.* 2008, Bhaduria *et al.* 2007, de Groot *et al.* 2007, Gasser *et al.* 2007, Gowda *et al.* 2006, Hauser *et al.* 2009, Kasuga *et al.* 2005, Lashkari *et al.* 1997, Mogensen *et al.* 2006, Rossouw *et al.* 2008). However, with the development of RNA (cDNA) sequencing, transcriptomics

© 2011 International Mycological Association


You are free to share - to copy, distribute and transmit the work, under the following conditions:

Attribution: You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).

Non-commercial: You may not use this work for commercial purposes.

No derivative works: You may not alter, transform, or build upon this work.

For any reuse or distribution, you must make clear to others the license terms of this work, which can be found at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>. Any of the above conditions can be waived if you get permission from the copyright holder. Nothing in this license impairs or restricts the author's moral rights.



GH family (Glycoside Hydrolase)	16
α -arabionofuranosidase	4
β -galactosidase	3
α -galactosidase	1
α -rhamnosidase	1
α -amylase	1
others	6
Metabolism	12
Transporter	7
Unknown	20

Fig. 1. Transcriptomics of *Aspergillus oryzae* during growth on L-arabinose. Left panel: Comparison of genes that are induced on L-arabinose to genes that are dependent on AraR. Right panel: functional annotation of the 55 genes that are induced by L-arabinose and regulated by AraR.

has become available for any species that has a sequenced genome and transcriptomic studies are now often included in genome papers. Proteomics was soon to follow and many studies on fungal proteomics have been reported in the last eight years (Acero *et al.* 2011, Grinyer *et al.* 2004, Ho *et al.* 2002, Kim *et al.* 2003, 2004, Lim *et al.* 2001, Matis *et al.* 2005).

These post-genomic studies have been aimed at many aspects of fungal biology. In this paper we present examples aimed at understanding how fungi interact with their environment. They were based on presentations at a Special Interest Group meeting during IMC9 in Edinburgh in August 2010.

CONTRIBUTIONS

Transcriptional regulation of genes involved in hemicellulose and cellulose utilization in *Aspergillus oryzae*

XlnR is a fungal transcription factor that regulates xylanolytic and cellulolytic enzymes in *Aspergillus*. *Aspergillus oryzae* XlnR was phosphorylated at low level in the absence of D-xylose, and was hyper-phosphorylated in its presence. Removal of D-xylose caused dephosphorylation of the hyper-phosphorylated forms leading to accumulation of less phosphorylated forms. XlnR activity is probably regulated by reversible phosphorylation (Noguchi *et al.* 2011).

While *Aspergillus niger* and *Aspergillus nidulans* have two XlnR homologs, AraR and GalA, *A. oryzae* has only AraR involved in L-arabinose catabolism. DNA microarray analysis for determining XlnR targets was performed by expression profiles of the XlnR overproducer and the XlnR deletion mutant after exposure to D-xylose for 30 min. The analysis revealed 75 genes as the possible targets of XlnR, including 32 glycoside hydrolases and three esterases for degradation of xylan and cellulose, seven transporters, and three genes for D-xylose catabolism (Noguchi *et al.* 2009). For AraR, two types of DNA microarray analysis were performed. One

compared expression profiles of L-arabinose induced and un-induced conditions, and the other compared those in the wild type and the AraR deletion mutant under L-arabinose induced conditions. By combination of the results, 55 genes were identified as possible AraR targets (Fig. 1), including 16 glycoside hydrolases, seven transporters, and several candidate genes for L-arabinose catabolism. The candidate genes for L-arabinose catabolism were expressed as His-tagged proteins in *Escherichia coli*. Analysis of substrate specificity of the recombinant enzymes led to identification of L-arabinose reductase and L-xylulose reductase in *A. oryzae*. The DNA microarray analysis contributed substantially to an in depth understanding of the degradation and utilization of hemicellulose by *A. oryzae*.

Zonal differentiation in sugar beet grown colonies of *Aspergillus niger*

Fungal colonies do not behave as uniform entities, but show differentiation for many aspects of physiology. The first report of this phenomenon described that protein secretion and growth only took place the periphery of colonies of *Aspergillus niger* grown on maltose (Wösten *et al.* 1991). To study this phenomenon in more detail, a specific growth system was developed called the ring-plate system (Levin *et al.* 2007b). This system consists of a round polycarbonate plate with concentric channels that are filled with liquid medium and a perforated polycarbonate membrane (Fig. 2). This membrane allows transport of nutrients and proteins, but the pores are too small to allow passage of fungal hyphae. Using this system it was shown that not only protein secretion and growth, but also overall gene expression is highly differentiated in colonies of *A. niger* grown on maltose or D-xylose (Levin *et al.* 2007a). Part of this differentiation could be attributed to depletion of the carbon source in the zones towards the centre of the colony. However, this was not the case for all gene systems. For instance, genes involved in nitrate utilisation were only expressed in the periphery of the colony, even though there was no significant difference in the nitrate concentration in the zones (Levin *et al.* 2007a).




Fig. 2. Schematic presentation of the ring-plate system . Reproduced with permission from Levin *et al.* (2007b).

In nature, fungi do not grow on high levels of pure mono- or disaccharides, but rather on mixtures of polymeric compounds (e.g. plant biomass). To release the monomeric carbon sources that can be taken up by the fungal cell, fungi produce complex mixtures of extracellular enzymes. The complexity of these mixtures was illustrated by a review presenting an overview of *Aspergillus* enzymes involved in plant cell wall degradation (de Vries & Visser 2001). This indicates that in nature fungi aren't subjected to a gradual reduction of a single carbon source, but more likely to substrates with changing compositions and levels.

To study zonal differentiation on natural substrates, *A. niger* was grown on the ring-plate system using sugar beet pulp as a carbon source. Sugar beet pulp is a waste product of the sugar industry and consists mainly of cellulose, xyloglucan and pectin. Growth on this substrate requires enzymatic hydrolysis of the polysaccharides.

Transcriptomic analysis using whole genome micro arrays for *A. niger* demonstrated that differentiation occurred to a lesser extent than on D-xylose and maltose (Benoit & de Vries, unpubl.). In addition, growth and protein secretion were observed throughout the colony, rather than only at the periphery (Benoit & de Vries, unpubl.).

Secretomes: Proteomic clues to fungal life-style choice

Many filamentous fungi have evolved to fill saprobic niches. This life-style, breaking down and utilizing complex biopolymers, requires a variety of hydrolytic enzymes secreted into the immediate environment of the fungus. Examination of the variations in the secretome of a fungus when confronted with different complex or simple substrates can reveal what the fungus is capable of utilizing and which secreted enzymes in its genome are expressed under particular conditions. We grew the zygomycete *Phycomyces blakesleeanus* and the ascomycete *Thielavia terrestris* on four complex carbon sources and glucose as a control using a defined medium of inorganic nutrients as a base. The complex

carbon sources were alder sawdust, pine sawdust, wheat bran and soybean hulls representing hardwoods, softwoods, grasses and dicotyledonous herbaceous plants respectively. The cultures were incubated at 30 °C (*P. blakesleeanus*) or 40 °C (*T. terrestris*) for four days and the fungal and plant solid material was removed by filtration through miracloth. The resulting liquid constituted the secretomes of the fungi. The proteins were concentrated on 10 kDa membranes, reduced and carboxamidomethylated, then digested with trypsin. The peptide solution was cleaned on disposable reverse phase (C18) columns and analyzed by high throughput liquid chromatography mass spectrometry (LC-MS) proteomics. The resulting MS and MS/MS spectra (molecular ion and fragmentation patterns, respectively) were analyzed against the protein databases of these two fungi using the SEQUEST program (Yates *et al.* 1998). The proteomics results for the zygomycete *P. blakesleeanus* suggest a bias towards scavenging protein with many proteases expressed under the different conditions (Table 1), but relatively few glycoside hydrolases. These proteomics observations are consistent with the paucity of glycoside hydrolase genes in this zygomycete's genome relative to saprobic ascomycetes. The ascomycete *T. terrestris* expressed a wide array of glycoside hydrolases, and some proteases, on the various substrates. The cellulose related CAZymes are shown as an example (Table 2). These proteomics results are consistent with the rich diversity of CAZymes found in saprobic ascomycete genomes (e.g. Martinez *et al.* 2008). Proteomics is a powerful technique for investigating which proteins are actually expressed and secreted by fungi when they are presented with different substrates in their environment.

Transcriptome profiling of the *Ustilago maydis* – maize interaction

Infection of maize by the fungal biotroph *Ustilago maydis* leads to formation of tumors in basically all aerial parts of the plant. Transcriptome profiling of *U. maydis* during pathogenic development was performed using a custom Affymetrix Gene-

Table 1. Proteases found in the secretomes of *Phycomyces blakesleeanus*. The Rhizopuspepsins represent a large family of A1A type proteases found in zygomycetes with six genes in *P. blakesleeanus*.

Glucose	Alder Sawdust	Pine Sawdust	Soybean Hulls	Wheat Bran	Hit Description	MEROPS Family
51	13	28	26	189	Rhizopuspepsin	A1A
		1	1	26	Rhizopuspepsin	A1A
2		4	29	20	Rhizopuspepsin	A1A
				15	Serine carboxypeptidase	S10
				14	Rhizopuspepsin	A1A
5					Aspartic protease	A1A
			5		Rhizopuspepsin	A1A

Table 2. CAZymes related to cellulose degradation found in the secretomes of *Thielavia terrestris* on various biomass sources (note that cellobiose dehydrogenase is not a glycoside hydrolase but is included as an enzyme important in the breakdown of cellulose).

Glucose	Alder Sawdust	Pine Sawdust	Soybean Hulls	Wheat Bran	Hit Description	GH Family
0	0	0	2	30	β -glucosidase	3
11	0	0	0	0	β -glucosidase	3
0	27	4	34	74	Endoglucanase	5
0	22	0	0	7	Endoglucanase	5
0	0	0	7	0	Endoglucanase	5
0	5	0	0	0	Endoglucanase	5
0	4	2	12	23	Exoglucanase	6
0	19	0	0	0	Exoglucanase	6
0	18	0	70	34	Endoglucanase	7
0	2	0	30	0	Exoglucanase	7
0	0	0	8	5	Exoglucanase	7
0	23	0	1	0	Exoglucanase	7
0	12	6	9	3	Endoglucanase	45
0	2	0	9	1	Endoglucanase	?
0	4	0	0	0	Cellobiose dehydrogenase	na

chip® microarray. This approach, together with sequencing of the *U. maydis* genome, identified 12 gene clusters encoding secreted effectors that are transcriptionally upregulated specifically during biotrophic interaction (Kämper *et al.* 2006). Deletion mutants for five of these gene clusters were altered in tumor formation, demonstrating the potential of transcript profiling for identifying virulence factors (Kämper *et al.* 2006)

A major step in understanding the impact of *U. maydis* infection on maize gene expression was achieved by transcriptome profiling of maize seedling leaves at different stages of infection (Doehlemann *et al.* 2008). Use of the Affymetrix maize genome array® identified a broad reprogramming of the maize primary- and secondary metabolism, particularly modulation of hormone signaling pathways as well as a shutdown of photosynthesis (Doehlemann *et al.* 2008). A major finding of this study was the attenuation of plant defences as soon as biotrophy has been established (Doehlemann *et al.* 2008).

Obviously there are fundamental differences between the various maize organs that *U. maydis* transforms into tumors. Therefore, transcriptome profiling was applied to study whether the distinctive developmental changes necessary for converting maize primordia to tumors would require organ-specific gene expression. To enable simultaneous transcript profiling of host and pathogen, a two organism microarray was designed using the Agilent® platform (Skibbe *et al.* 2010). Expression data from infected seedling leaf, adult leaf, and tassel revealed organ-specific expression programs of both interaction partners. In particular *U. maydis* genes encoding secreted effector proteins appeared to underlie organ-specific regulation (Skibbe *et al.* 2010). Moreover, *U. maydis* mutants deleted for clusters of secreted effectors (Kämper *et al.* 2006) showed significant differences in virulence depending on the infected maize organ (Skibbe *et al.* 2010). Together, these results show that tumor formation requires organ-specific gene expression by both partners. This finding

of organ-specific activity of pathogen effectors set a new paradigm in plant pathology and further demonstrates the power of transcriptome profiling in understanding of complex organismic interactions.

REFERENCES

- Acero FJ, Carbu M, El-Akhal MR, Garrido C, Gonzalez-Rodriguez VE, Cantoral JM (2011) Development of proteomics-based fungicides: new strategies for environmentally friendly control of fungal plant diseases. *International Journal of Molecular Sciences* **12**: 795–816.
- Andersen MR, Vongsangnak W, Panagiotou G, Salazar MP, Lehmann L, Nielsen J (2008) A trispecies *Aspergillus* microarray: comparative transcriptomics of three *Aspergillus* species. *Proceedings of the National Academy of Sciences, USA* **105**: 4387–4392.
- Bhadauria V, Popescu L, Zhao WS, Peng YL (2007) Fungal transcriptomics. *Microbiology Research* **162**: 285–298.
- Cuomo CA, Guldener U, Xu JR, Trail F, Turgeon BG, Di Pietro A, et al. (2007) The *Fusarium graminearum* genome reveals a link between localized polymorphism and pathogen specialization. *Science* **317**: 1400–1402.
- de Groot MJ, Daran-Lapujade P, van Breukelen B, Knijnenburg TA, de Hulster EA, Reinders MJ, et al. (2007) Quantitative proteomics and transcriptomics of anaerobic and aerobic yeast cultures reveals post-transcriptional regulation of key cellular processes. *Microbiology* **153**: 3864–3878.
- de Vries RP and Visser J (2001) *Aspergillus* enzymes involved in degradation of plant cell wall polysaccharides. *Microbiology and Molecular Biology Reviews* **65**: 497–522.
- Dean RA, Talbot NJ, Ebbole DJ, Farman ML, Mitchell TK, Orbach MJ, et al. (2005) The genome sequence of the rice blast fungus *Magnaporthe grisea*. *Nature* **434**: 980–986.
- Doehlemann G, Wahl R, Horst RJ, Voll LM, Usadel B, Poree F, et al. (2008) Reprogramming a maize plant: transcriptional and metabolic changes induced by the fungal biotroph *Ustilago maydis*. *Plant Journal* **56**: 181–195.
- Espagne E, Lespinet O, Malagnac F, Da Silva C, Jaillon O, Porcel BM, et al. (2008) The genome sequence of the model ascomycete fungus *Podospira anserina*. *Genome Biology* **9**: R77 (71–22).
- Galagan JE, Calvo SE, Borkovich KA, Selker EU, Read ND, Jaffe D, et al. (2003) The genome sequence of the filamentous fungus *Neurospora crassa*. *Nature* **422**: 859–868.
- Galagan JE, Calvo SE, Cuomo C, Ma LJ, Wortman JR, Batzoglou S, et al. (2005) Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* **438**: 1105–1115.
- Gasser B, Sauer M, Maurer M, Stadlmayr G, Mattanovich D (2007) Transcriptomics-based identification of novel factors enhancing heterologous protein secretion in yeasts. *Applied and Environmental Microbiology* **73**: 6499–6507.
- Gowda M, Venu RC, Raghupathy MB, Nobuta K, Li H, Wing R, et al. (2006) Deep and comparative analysis of the mycelium and appressorium transcriptomes of *Magnaporthe grisea* using MPSS, RL-SAGE, and oligoarray methods. *BMC Genomics* **7**: 310.
- Grinyer J, McKay M, Nevalainen H, Herbert BR (2004) Fungal proteomics: initial mapping of biological control strain *Trichoderma harzianum*. *Current Genetics* **45**: 163–169.
- Hauser NC, Dukalska M, Fellenberg K, Rupp S (2009) From experimental setup to data analysis in transcriptomics: copper metabolism in the human pathogen *Candida albicans*. *Journal of Biophotonics* **2**: 262–268.
- Ho Y, Gruhler A, Heilbut A, Bader GD, Moore L, Adams SL, et al. (2002) Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature* **415**: 180–183.
- Jeffries TW, Grigoriev IV, Grimwood J, Laplaza JM, Aerts A, Salamov A, et al. (2007) Genome sequence of the lignocellulose-bioconverting and xylose-fermenting yeast *Pichia stipitis*. *Nature Biotechnology* **25**: 319–326.
- Kämper J, Kahmann R, Bölker M, Ma L-J, Brefort T, Saville BJ, et al. (2006) Insights from the genome of the biotrophic fungal plant pathogen *Ustilago maydis*. *Nature* **444**: 97–101.
- Kasuga T, Townsend JP, Tian C, Gilbert LB, Mannhaupt G, Taylor JW, et al. (2005) Long-oligomer microarray profiling in *Neurospora crassa* reveals the transcriptional program underlying biochemical and physiological events of conidial germination. *Nucleic Acids Research* **33**: 6469–6485.
- Kim ST, Cho KS, Yu S, Kim SG, Hong JC, Han CD, et al. (2003) Proteomic analysis of differentially expressed proteins induced by rice blast fungus and elicitor in suspension-cultured rice cells. *Proteomics* **3**: 2368–2378.
- Kim ST, Yu S, Kim SG, Kim HJ, Kang SY, Hwang DH, et al. (2004) Proteome analysis of rice blast fungus (*Magnaporthe grisea*) proteome during appressorium formation. *Proteomics* **4**: 3579–3587.
- Lashkari DA, DeRisi JL, McCusker JH, Namath AF, Gentile C, Hwang SY, et al. (1997) Yeast microarrays for genome wide parallel genetic and gene expression analysis. *Proceedings of the National Academy of Sciences, USA* **94**: 13057–13062.
- Levin AM, de Vries RP, Conesa A, de Bekker C, Talon M, Menke HH, et al. (2007a) Spatial differentiation in the vegetative mycelium of *Aspergillus niger*. *Eukaryotic Cell* **6**: 2311–2322.
- Levin AM, de Vries RP and Wosten HA (2007b) Localization of protein secretion in fungal colonies using a novel culturing technique; the ring-plate system. *Journal of Microbiological Methods* **69**: 399–401.
- Lim D, Hains P, Walsh B, Bergquist P, Nevalainen H (2001) Proteins associated with the cell envelope of *Trichoderma reesei*: a proteomic approach. *Proteomics* **1**: 899–909.
- Machida M, Asai K, Sano M, Tanaka T, Kumagai T, Terai G, et al. (2005) Genome sequencing and analysis of *Aspergillus oryzae*. *Nature* **438**: 1157–1161.
- Martin F, Aerts A, Ahren D, Brun A, Danchin EG, Duchaussoy F, et al. (2008) The genome of *Laccaria bicolor* provides insights into mycorrhizal symbiosis. *Nature* **452**: 88–92.
- Martin F, Kohler A, Murat C, Balestrini R, Coutinho PM, Jaillon O, et al. (2010) Perigord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* **464**: 1033–1038.
- Martinez D, Berka RM, Henrissat B, Saloheimo M, Arvas M, Baker SE, et al. (2008) Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma reesei* (syn. *Hypocrea jecorina*). *Nature Biotechnology* **26**: 553–560.

- Martinez D, Larrondo LF, Putnam N, Gelpke MD, Huang K, Chapman J, *et al.* (2004) Genome sequence of the lignocellulose degrading fungus *Phanerochaete chrysosporium* strain RP78. *Nature Biotechnology* **22**: 695–700.
- Matis M, Zakej-Mavric M, Peter-Katalinic J (2005) Mass spectrometry and database search in the analysis of proteins from the fungus *Pleurotus ostreatus*. *Proteomics* **5**: 67–75.
- Mogensen J, Nielsen HB, Hofmann G, Nielsen J (2006) Transcription analysis using high-density micro-arrays of *Aspergillus nidulans* wild-type and *creA* mutant during growth on glucose or ethanol. *Fungal Genetics and Biology* **43**: 593–603.
- Nierman WC, Pain A, Anderson MJ, Wortman JR, Kim HS, Arroyo J, *et al.* (2005) Genomic sequence of the pathogenic and allergenic filamentous fungus *Aspergillus fumigatus*. *Nature* **438**: 1151–1156.
- Noguchi Y, Sano M, Kanamaru K, Ko T, Takeuchi M, Kato M, *et al.* (2009) Genes regulated by AoXlnR, the xylanolytic and cellulolytic transcriptional regulator, in *Aspergillus oryzae*. *Applied Microbiology and Biotechnology* **85**: 141–154.
- Ohm RA, de Jong JF, Lugones LG, Aerts A, Kothe E, Stajich JE, *et al.* (2010) Genome sequence of the model mushroom *Schizophyllum commune*. *Nature Biotechnology*. **28**: 957–963.
- Pel HJ, de Winde JH, Archer DB, Dyer PS, Hofmann G, Schaap PJ, *et al.* (2007) Genome sequencing and analysis of the versatile cell factory *Aspergillus niger* CBS 513.88. *Nature Biotechnology* **25**: 221–231.
- Rossouw D, Naes T, Bauer FF (2008) Linking gene regulation and the exo-metabolome: a comparative transcriptomics approach to identify genes that impact on the production of volatile aroma compounds in yeast. *BMC Genomics* **9**: 530.
- Skibbe DS, Doehlemann G, Fernandes J, Walbot V (2010) Maize tumors caused by *Ustilago maydis* require organ-specific genes in host and pathogen. *Science* **328**: 89–92.
- Wösten HAB, Mouhka SM, McLaughlin PMJ, Sietsma JH, Wessels JGH (1991) Localization of growth and excretion of proteins in *Aspergillus niger*. *Journal of General Microbiology* **137**: 2017–2023.
- Yates JR, Morgan SF, Gatlin CL, Griffin PR, Eng JK (1998) Method to compare collision-induced dissociation spectra of peptides: potential for library searching and subtractive analysis. *Analytical Chemistry* **70**: 3557–3565.